



ARMALA



Malathion 50% EC

Is a pesticide that is widely used in agriculture, residential landscaping, public recreation areas, and in public health pest control programs such as mosquito eradication:

- Mosquitoes (for eradication in public health, agriculture, and horticulture)
- Bollworms
- Mites
- Mustard Aphids
- Head Borers
- Tobacco Caterpillars

Always read and follow label instructions.

ARMALA: TOTAL PEST CONTROL FOR HEALTHIER CROPS AND SAFER ENVIRONMENTS

Active Ingredient:

Malathion 50% EC

Features:

- It provides a knockdown effect
- Used for control of several pests of small grains
- Used for direct contact sprays against sucking insects or as residual sprays against chewing pests

Mode of Action:

It kills pests by contact and ingestion.

Recommended Dose:

1-1.5 ml /1 L of water

Always read and follow label instructions.

 info@arbudaagrochemicals.com

Antidote:

- Atropinize the patient immediately and maintain full atropinization by a repeated dose of 2 to 4mg of atropine sulphate intravenously at 5 to 10 minutes intervals. As much as 25 to 50 mg of atropine may be required in a day. The need for further atropine administration is guided by the continuance of symptoms. The extent of salivation is a useful criterion for dose adjustment.
- Dissolve 1-2 gm 2-PAM in 10 ml distilled water and inject slowly IV over 10-15 minutes.

Packing:

1 L, 5 L, 20 L

Scan QR CODE
for product details



Always read and follow label instructions.

 info@arbudaagrochemicals.com

 +91 9076 9076 42

 www.arbudaagrochemicals.com